

A QUANTITATIVE ANALYSIS OF METRICS ON \mathbb{R}^n WITH ALMOST CONSTANT POSITIVE SCALAR CURVATURE, WITH APPLICATIONS TO FAST DIFFUSION FLOWS

G. CIRAOLLO, A. FIGALLI, AND F. MAGGI

ABSTRACT. We prove a quantitative structure theorem for metrics on \mathbb{R}^n that are conformal to the flat metric, have almost constant positive scalar curvature, and cannot concentrate more than one bubble. As an application of our result, we show a quantitative rate of convergence in relative entropy for a fast diffusion equation in \mathbb{R}^n related to the Yamabe flow.

1. INTRODUCTION

1.1. The prescribed scalar curvature problem on the sphere. Given a n -dimensional Riemannian manifold (M, g_0) , $n \geq 3$, the problem of finding a metric g conformal to g_0 whose scalar curvature R_g is equal to a prescribed function R boils down to showing the existence of a positive solution u to the nonlinear PDE

$$-\Delta_{g_0} u + \frac{n-2}{4(n-1)} R_{g_0} u = \frac{n-2}{4(n-1)} R u^p \quad (1.1)$$

where $\Delta_{g_0} = \operatorname{div}(\nabla_{g_0} \cdot)$ and R_{g_0} denote respectively the Laplace-Beltrami operator and the scalar curvature of (M, g_0) , and

$$p = \frac{n+2}{n-2} = 2^* - 1, \quad 2^* = \frac{2n}{n-2}.$$

Indeed, if u solves (1.1), then the metric $g = u^{p-1} g_0$ satisfies $R_g = R$.

When (M, g_0) is the round sphere then $R_{g_0} = n(n-1)$ and (1.1) can be read on \mathbb{R}^n by means of the stereographic projection. More precisely, consider the inverse stereographic projection $F : \mathbb{R}^n \rightarrow \mathbb{S}^n$ defined by

$$F(x) = \left(\frac{2x}{1+|x|^2}, \frac{|x|^2-1}{1+|x|^2} \right). \quad (1.2)$$

Then $v : \mathbb{S}^n \rightarrow \mathbb{R}$ solves (1.1) if and only if

$$u(x) = \left(\frac{2}{1+|x|^2} \right)^{(n-2)/2} v(F(x))$$

solves

$$-\Delta u = K u^p \quad \text{on } \mathbb{R}^n, \quad (1.3)$$

where $K(x) = \frac{n-2}{4(n-1)} R(F(x))$.

When looking for solutions of (1.3), it is natural to impose that u satisfies

$$u > 0 \text{ on } \mathbb{R}^n \quad \text{and} \quad \int_{\mathbb{R}^n} |\nabla u|^2 < \infty. \quad (1.4)$$

In general, for a given function K (and even when K is just a small perturbation of a constant) there may exist no solution to (1.3)–(1.4) (see [KW74]), and indeed there is a vast literature dedicated to finding necessary and sufficient conditions on K in order to guarantee the solvability of (1.3)–(1.4), see for example [Mos73, Ni82, DN85, ES86, Che89, CY91, BC91, BE93, Li95, Li96, Bah96, CL97, AGAP99, AM01, LL09, LZ15]. At the same time, when K is constantly equal to

some $\kappa > 0$, the problem is completely rigid. Indeed, by [Oba72, GNN79], if u solves (1.3)–(1.4) with $K \equiv \kappa > 0$, then there exist $\lambda > 0$ and $z \in \mathbb{R}^n$ such that

$$u(x) = \lambda^{(n-2)/2} v_\kappa(\lambda(x - z)) \quad \forall x \in \mathbb{R}^n,$$

where

$$v_\kappa(x) = \left(\frac{n(n-2)}{\kappa} \right)^{(n-2)/4} \frac{1}{(1 + |x|^2)^{(n-2)/2}} \quad \forall x \in \mathbb{R}^n. \quad (1.5)$$

1.2. Main result. The goal of this paper is to give a quantitative description of solutions u to the *prescribed scalar curvature equation* (1.3)–(1.4) in the regime when K is close (in a suitable sense) to a positive constant.

In order to identify the natural space in which the distance of K to a constant should be measured, we make the following observation:

$$-\Delta u = K u^p \quad \Longleftrightarrow \quad \int_{\mathbb{R}^n} \nabla u \cdot \nabla \varphi = \int_{\mathbb{R}^n} K u^p \varphi \quad \forall \varphi \in \dot{W}^{1,2}(\mathbb{R}^n),$$

where $\dot{W}^{1,2}(\mathbb{R}^n)$ denotes the closure of $C_c^\infty(\mathbb{R}^n)$ with respect to the norm $\|\nabla \cdot\|_{L^2(\mathbb{R}^n)}$. Since

$$\varphi \in \dot{W}^{1,2}(\mathbb{R}^n) \hookrightarrow L^{2^*}(\mathbb{R}^n),$$

$\int_{\mathbb{R}^n} K u^p \varphi$ is well-defined provided $K u^p$ belongs to the dual of $L^{2^*}(\mathbb{R}^n)$, namely $L^{2n/(n+2)}(\mathbb{R}^n)$.

This suggests the following definition:

$$\delta(u) := \|K u^p - K_0(u) u^p\|_{L^{2n/(n+2)}} = \left(\int_{\mathbb{R}^n} \left| \frac{\Delta u}{u^p} + K_0(u) \right|^{2n/(n+2)} u^{2^*} \right)^{(n+2)/2n}, \quad (1.6)$$

where

$$K_0(u) := \frac{\int_{\mathbb{R}^n} K u^{2^*}}{\int_{\mathbb{R}^n} u^{2^*}} = \frac{\int_{\mathbb{R}^n} \frac{-\Delta u}{u^p} u^{2^*}}{\int_{\mathbb{R}^n} u^{2^*}} = \frac{\int_{\mathbb{R}^n} |\nabla u|^2}{\int_{\mathbb{R}^n} u^{2^*}}. \quad (1.7)$$

Hence, the question becomes: if $\delta(u)$ is small, can we say that u is close to a translation/dilation of $v_{K_0(u)}$?

A negative answer is given by the following simple example: given a function $v : \mathbb{R}^n \rightarrow \mathbb{R}$, set

$$v[z, \lambda](x) := \lambda^{(n-2)/2} v(\lambda(x - z)), \quad x \in \mathbb{R}^n,$$

and consider

$$u = \sum_{i=1}^m v_1[z_i, \lambda_i],$$

where the functions $v_1[z_i, \lambda_i]$ are supported far away from each other. Then

$$-\Delta u = -\sum_{i=1}^m \Delta v_1[z_i, \lambda_i] = \sum_{i=1}^m v_1[z_i, \lambda_i]^p = K \left(\sum_{i=1}^m v_1[z_i, \lambda_i] \right)^p = K u^p,$$

with

$$K = \frac{\sum_{i=1}^m v_1[z_i, \lambda_i]^p}{\left(\sum_{i=1}^m v_1[z_i, \lambda_i] \right)^p},$$

and it is easy to check that by taking the point z_i sufficiently far from each other one can make $\|K u^p - u^p\|_{L^{2n/(n+2)}}$ arbitrarily small.

As shown by Struwe's [Str84] (see also [Heb14, Theorem 3.3]), this bubbling phenomenon is the only possible “bad” case. More precisely, whenever $\delta(u)$ is small, u is close to a sum of bubbles as above. Luckily, in many applications, this phenomenon can be avoided by some preliminary study of the PDE under investigation, and one can usually localize the problem in a suitable way so that, in the region under investigation, the “energy” of u is strictly less than the energy of two bubbles. Hence, we shall focus on the latter situation.

Before stating our result, we recall the definition of the Sobolev constant on \mathbb{R}^n ,

$$S = \inf \left\{ \frac{\|\nabla v\|_{L^2(\mathbb{R}^n)}}{\|v\|_{L^{2^*}(\mathbb{R}^n)}} : v \neq 0, |\{|v| > t\}| < \infty \quad \forall t > 0 \right\}. \quad (1.8)$$

By [Aub76, Tal76, CENV04], the family of functions $\{v_\kappa[z, \lambda]\}_{\kappa, \lambda, z}$ corresponds to the minimizers in (1.8). Hence, since $-\Delta v_\kappa = \kappa v_\kappa^p$, one can easily check that

$$\int_{\mathbb{R}^n} |\nabla v_\kappa|^2 = S^2 \left(\int_{\mathbb{R}^n} v_\kappa^{2^*} \right)^{2/2^*} = \frac{S^n}{\kappa^{(n-2)/2}} \quad \int_{\mathbb{R}^n} v_\kappa^{2^*} = \frac{S^n}{\kappa^{n/2}}. \quad (1.9)$$

To simplify the notation (and because for most applications this is not a real restriction), we shall assume that the energy of u is bounded by $3/2$ the energy of a single bubble. Of course $3/2$ does not play any essential role, and the proof holds when $3/2$ is replaced by any constant strictly less than 2. Also, the assumption $K_0(u) = 1$ is not restrictive, since it can always be guaranteed by rescaling u .

Theorem 1.1. *Given $n \geq 3$, there exists $C_0 = C_0(n) > 0$ with the following property. Let $u \in C^\infty(\mathbb{R}^n) \cap \dot{H}^1(\mathbb{R}^n)$ be a positive function satisfying*

$$K_0(u) = 1 \quad \text{and} \quad \int_{\mathbb{R}^n} |\nabla u|^2 \leq \frac{3}{2} S^n.$$

Then there exist $z \in \mathbb{R}^n$ and $\lambda \in (0, \infty)$ such that

$$u = v_1[z, \lambda] + \rho,$$

where

$$\|\nabla \rho\|_{L^2(\mathbb{R}^n)} \leq C_0 \delta(u). \quad (1.10)$$

Remark 1.2. Theorem 1.1 is easily seen to be optimal. Indeed, set $v := v_1[0, 1]$, let $\varepsilon > 0$ be small, and consider $u_\varepsilon := v_1 + \varepsilon \phi$ for some $\phi \in C_c^\infty(\mathbb{R}^n)$, with $\phi \geq 0$. Then

$$\Delta u_\varepsilon = \Delta v + \varepsilon \Delta \phi = v^p + \varepsilon \Delta \phi = K_\varepsilon u_\varepsilon^p$$

with

$$K_\varepsilon := \frac{v^p + \varepsilon \Delta \phi}{u_\varepsilon^p} = 1 + \frac{[v^p - (v + \varepsilon \phi)^p] + \varepsilon \Delta \phi}{u_\varepsilon^p} = 1 + O(\varepsilon u_\varepsilon^{-p}).$$

This shows that

$$\|K_\varepsilon u_\varepsilon^p - u_\varepsilon^p\|_{L^{2n/(n+2)}} = O(\varepsilon). \quad (1.11)$$

Also, since

$$K_0(u_\varepsilon) = \frac{\int_{\mathbb{R}^n} K_\varepsilon u_\varepsilon^{2^*}}{\int_{\mathbb{R}^n} u_\varepsilon^{2^*}},$$

by (1.11) and Hölder inequality,

$$|K_0(u_\varepsilon) - 1| \leq \frac{|\int_{\mathbb{R}^n} (K_\varepsilon - 1) u_\varepsilon^{2^*}|}{\int_{\mathbb{R}^n} u_\varepsilon^{2^*}} \leq \left(\frac{\int_{\mathbb{R}^n} |K_\varepsilon - 1|^{2n/(n+2)} u_\varepsilon^{2^*}}{\int_{\mathbb{R}^n} u_\varepsilon^{2^*}} \right)^{(n+2)/2n} = O(\varepsilon).$$

Combining this bound with (1.11) we get $\delta(u_\varepsilon) \leq C\varepsilon \simeq \|\nabla(\varepsilon \phi)\|_{L^2(\mathbb{R}^n)}$, which proves the optimality of our result.

As an application of Theorem 1.1, we investigate the behavior of solutions to a fast diffusion equation in \mathbb{R}^n related to the Yamabe flow.

1.3. Convergence to equilibrium: a fast diffusion equation related to the Yamabe flow. Given $m \in (0, 1)$, the Cauchy problem for the fast diffusion equation is written as

$$\frac{d}{dt}u = \Delta(u^m) \quad \text{in } (0, \infty) \times \mathbb{R}^n. \quad (1.12)$$

Assuming that the initial datum u_0 is nonnegative and fastly decaying at infinity, it is well-known that solutions to (1.12) are smooth and positive for all times if $m > m_c = (n-2)/n$, while they vanish in finite time if $m \leq m_c$. There is a huge literature on the subject, and we refer the interested reader to the monograph [Váz06] for a comprehensive overview and more references.

A case of a special interest corresponds to the choice $m = (n-2)/(n+2)$, where the equation for u is equivalent to the (not volume-preserving) Yamabe flow

$$\frac{d}{dt}g = -R_g g$$

for the metric $g_{ij}(t) = u(t)^{p-1} dx_i dx_j$ (recall that $p = 2^* - 1$). Hence, we consider the Cauchy problem

$$\frac{d}{dt}u = \Delta(u^m) \quad \text{in } (0, \infty) \times \mathbb{R}^n, \quad m = \frac{n-2}{n+2} = \frac{1}{p}, \quad (1.13)$$

with a continuous initial datum $u_0 \geq 0$ satisfying $u_0(x) = O(|x|^{-(n+2)})$ as $|x| \rightarrow \infty$. Because of its geometric relevance, this equation has received a lot of attention. In particular, as shown in [dPS01, Theorem 1.1] (see also [Cho92, Ye94]), under the above assumptions on u_0 there exists a vanishing time $T = T(u_0) > 0$, a point $z \in \mathbb{R}^n$, and a number $\lambda > 0$, such that $u \equiv 0$ for $t \geq T$ and

$$\left\| \frac{(T-t)^{-1/(1-m)} u(t, \cdot)}{v_{1/(1-m)}[z, \lambda]^{1/m}} - 1 \right\|_{L^\infty(\mathbb{R}^n)} \rightarrow 0 \quad \text{as } t \rightarrow T^-. \quad (1.14)$$

Later on, in [DS08, BBD⁺09, BDGV10, BGV10], the authors investigated the asymptotic of solutions for all values of $m \in (0, 1)$, and proved both qualitative and quantitative convergence results under the assumption that the initial datum is trapped in between two Barenblatt solutions with the same extinction time.

As observed in [DS08] (see also [DKS] for a different but related analysis), this trapping assumption on the initial datum is very restrictive in our setting, as it completely misses the picture given by [dPS01] and gives rise to an extinction profile different from the one in (1.14), which is believed to be the correct one for “most” initial data. Indeed, as shown in [DS08, Theorem 1.4], there exists a large class of initial data of the form

$$u_0(x) = \frac{C_0}{|x|^{(n+2)/2}} (1 + o(1)) \quad \text{as } |x| \rightarrow \infty, \quad C_0 > 0,$$

that do not satisfy the trapping assumption and whose solutions behave as follows: there exists a time $t_0 \in (0, T)$ (which can be explicitly computed in terms of C_0 and is given by $t_0 = \frac{C_0(n+2)}{(n-2)^2}$) such that the behavior of the solution is governed by a Barenblatt profile $B(t, \cdot)$ up to the time t_0 when the Barenblatt vanishes. Then the solution develops a singularity at $t = t_0$, and it satisfies $u(t, x) = O(|x|^{-(n+2)})$ as $|x| \rightarrow \infty$ for all $t > t_0$ (see [DS08, Theorem 1.4(ii)]). In particular, as observed in [DS08, Corollary 1.5], this allows one to apply [dPS01, Theorem 1.1] and deduce that $u(t, x)$ exhibits the vanishing profile of a sphere, as shown by (1.14).

By exploiting our Theorem 1.1, we can improve the convergence result in (1.14) and obtain a quantitative rate of convergence under the same assumptions as in [dPS01]. More precisely, in section 3 we prove the following:

Theorem 1.3. *Let u be a solution to the fast diffusion equation (1.13) starting from a non-negative continuous initial datum u_0 satisfying $u_0(x) = O(|x|^{-(n+2)})$ as $|x| \rightarrow \infty$. Let $T =$*

$T(u_0) > 0$ denote the vanishing time of u . Then there exist $z \in \mathbb{R}^n$, $\lambda > 0$, and a dimensional constant $\kappa(n) > 0$ such that

$$\left\| \frac{(T-t)^{-1/(1-m)} u(t, \cdot)}{v_{1/(1-m)}[z, \lambda]^{1/m}} - 1 \right\|_{L^\infty(\mathbb{R}^n)} \leq C_* (T-t)^{\kappa(n)} \quad \forall 0 < t < T,$$

where $C_* > 0$ is a constant depending on the initial datum u_0 .

Acknowledgment: The authors wish to thank Manuel Del Pino for suggesting the possibility of improving [dPS01, Theorem 1.1]. This work has been done while GC was visiting the University of Texas at Austin under the support of NSF FRG Grant DMS-1361122, the Oden Fellowship at ICES, the GNAMPA of the Istituto Nazionale di Alta Matematica (INdAM), and the FIR project 2013 “Geometrical and Qualitative aspects of PDE”. AF was partially supported by NSF Grants DMS-1262411 and DMS-1361122, and FM was supported by NSF Grants DMS-1265910 and DMS-1361122.

2. PROOF OF THEOREM 1.1

This section is devoted to the proof of Theorem 1.1. Recall the notation

$$n \geq 3, \quad 2^* = \frac{2n}{n-2}, \quad p = 2^* - 1 = \frac{n+2}{n-2} = 1 + s, \quad s = p - 1 = \frac{4}{n-2}.$$

We begin by observing that it is enough to prove the theorem for $u : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfying

$$u \in C^\infty(\mathbb{R}^n) \cap \dot{H}^1(\mathbb{R}^n), \quad u > 0 \text{ on } \mathbb{R}^n,$$

and such that

$$K_0(u) = 1, \quad \int_{\mathbb{R}^n} |\nabla u|^2 \leq \frac{3}{2} S^n, \quad \delta = \delta(u) \leq \delta_0, \quad (2.1)$$

for a suitably small constant $\delta_0 = \delta_0(n)$.

Indeed we note that if $\delta(u) > \delta_0$, then the theorem is trivially true simply by choosing $\lambda = 1$, $z = 0$, setting

$$\rho = u - v_1,$$

and then simply choosing $C_0 = C_0(n)$ large enough.

Step one. Thanks to [Str84], for a constant $\varepsilon_0 = \varepsilon_0(n) > 0$ to be fixed later on, we can choose δ_0 depending on ε_0 in such a way that there exist $z \in \mathbb{R}^n$, $\lambda, \alpha > 0$, such that

$$\begin{aligned} \left\| \nabla u - \alpha \nabla v_1[z, \lambda] \right\|_{L^2(\mathbb{R}^n)} &\leq \varepsilon_0, \\ |\alpha - 1| &\leq \varepsilon_0. \end{aligned} \quad (2.2)$$

Without loss of generality, the parameters $z \in \mathbb{R}^n$ and $\lambda, \alpha \in (0, \infty)$ can be chosen in such a way that

$$\left\| \nabla u - \alpha \nabla v_1[z, \lambda] \right\|_{L^2(\mathbb{R}^n)} = \min_{w \in \mathbb{R}^n, \mu, a > 0} \left\| \nabla u - a \nabla v_1[w, \mu] \right\|_{L^2(\mathbb{R}^n)}. \quad (2.3)$$

In particular, if we set

$$\begin{aligned} \rho &= u - \sigma, \quad \sigma = \alpha U, \\ U &= v_1[z, \lambda], \quad V = \frac{\partial v_1[w, \mu]}{\partial \mu} \Big|_{w=z, \mu=\lambda}, \quad W^j = \frac{\partial v_1[w, \mu]}{\partial w_j} \Big|_{w=z, \mu=\lambda}, \end{aligned}$$

then by (2.3) we find that

$$\int_{\mathbb{R}^n} \nabla U \cdot \nabla \rho = \int_{\mathbb{R}^n} \nabla V \cdot \nabla \rho = \int_{\mathbb{R}^n} \nabla W^j \cdot \nabla \rho = 0, \quad \forall 1 \leq j \leq n. \quad (2.4)$$

Also, the first bound in (2.2) gives

$$\|\nabla \rho\|_{L^2(\mathbb{R}^n)} \leq \varepsilon_0. \quad (2.5)$$

By a spectral analysis argument (see, e.g., the appendix to [BE91]), (2.4) implies that

$$\int_{\mathbb{R}^n} |\nabla \rho|^2 \geq \Lambda \int_{\mathbb{R}^n} U^{p-1} \rho^2 \quad (2.6)$$

where $\Lambda = \Lambda(n)$ is such that

$$\Lambda > p.$$

Step two. Having in mind to exploit $\Lambda > p$, we now test the equation $-\Delta u = K u^p$ with ρ , and using $\int_{\mathbb{R}^n} \nabla \rho \cdot \nabla U = 0$ we get

$$\int_{\mathbb{R}^n} |\nabla \rho|^2 = \int_{\mathbb{R}^n} K u^p \rho = \int_{\mathbb{R}^n} u^p \rho + \int_{\mathbb{R}^n} (K-1) u^p \rho. \quad (2.7)$$

Since $u = \sigma + \rho$, a Taylor expansion yields

$$\int_{\mathbb{R}^n} u^p \rho = \int_{\mathbb{R}^n} \sigma^p \rho + p \int_{\mathbb{R}^n} \sigma^{p-1} \rho^2 + O\left(\int_{\mathbb{R}^n} |\nabla \rho|^2\right)^{1+\gamma}, \quad (2.8)$$

where

$$\gamma = \min\left\{\frac{1}{2}, \frac{2}{n-2}\right\}.$$

From $\sigma^p = \alpha^p U^p = -\alpha^p \Delta U$ and using again that $\int_{\mathbb{R}^n} \nabla \rho \cdot \nabla U = 0$ we get

$$\int_{\mathbb{R}^n} |\nabla \rho|^2 - p \int_{\mathbb{R}^n} \sigma^{p-1} \rho^2 = \int_{\mathbb{R}^n} (K-1) u^p \rho + O\left(\int_{\mathbb{R}^n} |\nabla \rho|^2\right)^{1+\gamma}.$$

Note that, by Hölder inequality (recall that $p = 2^*/(2^*)'$ and $(2^*)' = 2n/(n+2)$)

$$\left| \int_{\mathbb{R}^n} (K-1) u^p \rho \right| \leq \delta(u) \|\rho\|_{L^{2^*}(\mathbb{R}^n)} \leq S \|\nabla \rho\|_{L^2(\mathbb{R}^n)} \delta(u). \quad (2.9)$$

Also, recalling (2.6),

$$\int_{\mathbb{R}^n} \sigma^{p-1} \rho^2 = \alpha^{p-1} \int_{\mathbb{R}^n} U^{p-1} \rho^2 \leq \frac{\alpha^{p-1}}{\Lambda} \int_{\mathbb{R}^n} |\nabla \rho|^2.$$

Hence

$$\left(1 - \frac{\alpha^{p-1} p}{\Lambda}\right) \int_{\mathbb{R}^n} |\nabla \rho|^2 \leq S \|\nabla \rho\|_{L^2(\mathbb{R}^n)} \delta(u) + O\left(\int_{\mathbb{R}^n} |\nabla \rho|^2\right)^{1+\gamma}.$$

Since $\Lambda > p$ and $|\alpha - 1| \leq \varepsilon_0$, choosing $\delta(u)$ small enough we can ensure that $1 - \frac{\alpha^{p-1} p}{\Lambda} \geq c_0 > 0$ for some dimensional constant c_0 , and we get

$$c_0 \int_{\mathbb{R}^n} |\nabla \rho|^2 \leq S \|\nabla \rho\|_{L^2(\mathbb{R}^n)} \delta(u) + O\left(\int_{\mathbb{R}^n} |\nabla \rho|^2\right)^{1+\gamma}.$$

Since $\int_{\mathbb{R}^n} |\nabla \rho|^2$ is smaller than ε_0 , we can also reabsorb the last term to conclude that $\|\nabla \rho\|_2 \leq C \delta(u)$.

Step three. We now quantitatively control $|\alpha - 1|$. To this aim, we observe that assumption $K_0(u) = 1$ is equivalent to

$$\int_{\mathbb{R}^n} |\nabla u|^2 = \int_{\mathbb{R}^n} u^{2^*}$$

(see (1.7)). Note that, since $\|\nabla \rho\|_2 \leq C \delta(u)$ (by Step 2),

$$\int_{\mathbb{R}^n} |\nabla u|^2 = \int_{\mathbb{R}^n} |\nabla \sigma|^2 + \int_{\mathbb{R}^n} |\nabla \rho|^2 = \alpha^2 S^n + O(\delta^2).$$

On the other hand, recalling that $\int_{\mathbb{R}^n} \sigma^p \rho = 0$ (cp. with Step two)

$$\int_{\mathbb{R}^n} u^{2^*} = \int_{\mathbb{R}^n} \sigma^{2^*} + p \int_{\mathbb{R}^n} \sigma^p \rho + O(\delta^2) = \alpha^{2^*} S^n + O(\delta^2).$$

Comparing these expressions, we immediately deduce that

$$\alpha^{2^*} S^n = \alpha^2 S^n + O(\delta^2),$$

hence $|\alpha - 1| \leq O(\delta^2)$. Thus, if we set $\rho' := \rho + (\alpha - 1)U$, we proved that

$$u = U + \rho'$$

where $\|\nabla \rho'\|_2 \leq C_0 \delta(u)$, as desired. \square

3. AN APPLICATION TO FAST DIFFUSION EQUATIONS: PROOF OF THEOREM 1.3

This section is devoted to the proof of Theorem 1.3. Thus, we consider a continuous initial datum $u_0 \geq 0$ satisfying $u_0(x) = O(|x|^{-(n+2)})$ as $|x| \rightarrow \infty$, and u a solution of (1.13) with $u(0, \cdot) = u_0$. As explained in Section 1.3, under these assumptions the qualitative convergence result (1.14) holds, and our goal is to quantify the rate of convergence.

We first notice that (1.14) can be restated as follows (see [dPS01, Theorem 1.1]): there exists a vanishing time $T > 0$ (depending on u_0) such that $u \equiv 0$ for $t \geq T$ and

$$\frac{u(t, x)}{(T - t)^{1/(1-m)}} = v_{1/(1-m)}[z_\infty, \lambda_\infty]^{1/m} + \theta(t, x) \quad \text{for } t < T, \quad (3.1)$$

where $\lambda_\infty > 0$, $z_\infty \in \mathbb{R}^n$, and θ satisfy

$$\sup_{x \in \mathbb{R}^n} (1 + |x|^{n+2}) |\theta(t, x)| \rightarrow 0 \quad \text{as } t \rightarrow T^-. \quad (3.2)$$

For proving Theorem 1.3 we need to show the existence of $\kappa(n) > 0$ such that

$$\sup_{x \in \mathbb{R}^n} (1 + |x|^{n+2}) |\theta(t, x)| \leq C(T - t)^{\kappa(n)} \quad \text{for } 0 < t < T. \quad (3.3)$$

Following [dPS01], define

$$w(s, x) = \frac{u(t, x)^m}{(T - t)^{m/(1-m)}} \Big|_{t=T(1-e^{-s})}, \quad W_\infty(x) = v_{1/(1-m)}[z_\infty, \lambda_\infty](x),$$

so that (3.1) and (3.2) imply, setting for short $w(s) = w(s, \cdot)$,

$$\lim_{s \rightarrow +\infty} \|w(s) - W_\infty\|_{L^{2^*}(\mathbb{R}^n)} = 0. \quad (3.4)$$

Recalling the notation $p = 2^* - 1$, we see that w satisfies

$$\frac{d}{ds} w^p = \Delta w + \frac{1}{1-m} w^p \quad \text{on } (0, \infty) \times \mathbb{R}^n, \quad (3.5)$$

while of course $W = W_\infty$ is a solution to

$$\Delta W + \frac{1}{1-m} W^p = 0 \quad \text{on } \mathbb{R}^n. \quad (3.6)$$

Let us consider the functional

$$J[v] = \int_{\mathbb{R}^n} \frac{|\nabla v|^2}{2} - \frac{1}{1-m} \int_{\mathbb{R}^n} \frac{v^{2^*}}{2^*}.$$

By (3.5) we compute

$$\frac{d}{ds} J[w(s)] = - \int_{\mathbb{R}^n} \left(\Delta w(s) + \frac{1}{1-m} w(s)^p \right) \frac{d}{ds} w(s) = - \frac{1}{p} \int_{\mathbb{R}^n} \left(\frac{\Delta w(s)}{w(s)^p} + \frac{1}{1-m} \right)^2 w(s)^{2^*}, \quad (3.7)$$

so that $s \mapsto J[w(s)]$ is decreasing, with $J[w(s)] \geq J[W_\infty]$ by Fatou's lemma and (3.4). Exploiting the fact that

$$\mathbb{R} \ni c \mapsto \int_{\mathbb{R}^n} \left(\frac{\Delta w(s)}{w(s)^p} - c \right)^2 w(s)^{2^*}$$

attains its minimum at

$$c = \frac{\int_{\mathbb{R}^n} \frac{\Delta w(s)}{w(s)^p} w(s)^{2^*}}{\int_{\mathbb{R}^n} w(s)^{2^*}} = - \frac{\int_{\mathbb{R}^n} |\nabla w(s)|^2}{\int_{\mathbb{R}^n} w(s)^{2^*}} = -K_0(w(s)),$$

by Hölder inequality we find that

$$\begin{aligned} \delta(w(s))^2 &\leq \left(\int_{\mathbb{R}^n} w(s)^{2^*} \right)^{2/n} \int_{\mathbb{R}^n} \left(\frac{\Delta w(s)}{w(s)^p} + K_0(w(s)) \right)^2 w(s)^{2^*} \\ &\leq C(n) \int_{\mathbb{R}^n} \left(\frac{\Delta w(s)}{w(s)^p} + \frac{1}{1-m} \right)^2 w(s)^{2^*}. \end{aligned} \quad (3.8)$$

Hence, thanks to (3.4), there exists s_0 (depending on the initial datum u_0) such that

$$c(n) \leq \int_{\mathbb{R}^n} w(s)^{2^*} \leq C(n) \quad \forall s \geq s_0. \quad (3.9)$$

Combining (3.7), (3.8), and (3.9) we find

$$\delta(w(s))^2 \leq -C(n) \frac{d}{ds} J[w(s)] \quad \forall s \geq s_0, \quad (3.10)$$

so that

$$\int_{s_0}^{\infty} \delta(w(s))^2 ds \leq C(n) (J[w(s_0)] - J[w(\infty)]) \leq C(n) (J[w(s_0)] - J[W_\infty]) < \infty.$$

In particular we can find a sequence $s_j \rightarrow \infty$ such that $\delta(w(s_j)) \rightarrow 0$ as $j \rightarrow \infty$. We can thus apply Struwe's theorem to $\{w(s_j)\}_{j \in \mathbb{N}}$. Since (3.4) excludes bubbling, we conclude that

$$\lim_{s \rightarrow \infty} J[w(s)] = \lim_{j \rightarrow \infty} J[w(s_j)] = J[W_\infty].$$

Furthermore, using again (3.4), this implies

$$\lim_{s \rightarrow \infty} \int_{\mathbb{R}^n} |\nabla w(s)|^2 = \int_{\mathbb{R}^n} |\nabla W_\infty|^2, \quad (3.11)$$

and thus also

$$\lim_{s \rightarrow \infty} \|\nabla w(s) - \nabla W_\infty\|_{L^2(\mathbb{R}^n)} = 0, \quad \lim_{s \rightarrow \infty} K_0(w(s)) = \frac{\int_{\mathbb{R}^n} |\nabla W_\infty|^2}{\int_{\mathbb{R}^n} W_\infty^{2^*}} = \frac{1}{1-m}. \quad (3.12)$$

Now, for any $s > 0$, denote by $W(s)$ the unique minimizer of

$$W \mapsto \|\nabla w(s) - \nabla W\|_{L^2(\mathbb{R}^n)} \quad (3.13)$$

among all positive functions W satisfying (3.6). Then, setting $w(s) = W(s) + \rho(s)$,

$$\int_{\mathbb{R}^n} \nabla W(s) \cdot \nabla \rho(s) = 0 = \int_{\mathbb{R}^n} W(s)^p \rho(s), \quad \lim_{s \rightarrow \infty} \|\nabla \rho(s)\|_{L^2(\mathbb{R}^n)} = 0, \quad (3.14)$$

where the last limit follow by (3.12) and by the minimality property of $W(s)$. Since J is constant on solutions of (3.6) (so $J[W(s)] = J[W_\infty]$), we can expand J around $W(s)$ with the aid of (3.14), to find out that

$$\begin{aligned} I[w(s)] &:= J[w(s)] - J[W_\infty] = J[w(s)] - J[W(s)] \\ &= \frac{1}{2} \int_{\mathbb{R}^n} |\nabla \rho(s)|^2 - \frac{1}{(1-m)2^*} \int_{\mathbb{R}^n} \left((W(s) + \rho(s))^{2^*} - W(s)^{2^*} - 2^* W(s)^p \rho(s) \right) \end{aligned}$$

$$\begin{aligned}
&\approx \int_{\mathbb{R}^n} |\nabla \rho(s)|^2 - \frac{p}{1-m} \int_{\mathbb{R}^n} W(s)^{p-1} \rho(s)^2 \\
&\approx \int_{\mathbb{R}^n} |\nabla \rho(s)|^2 = \int_{\mathbb{R}^n} |\nabla w(s) - \nabla W(s)|^2 \quad \forall s \geq s_0,
\end{aligned} \tag{3.15}$$

where $a \approx b$ means that $a/C(n) \leq b \leq C(n)a$ for some positive dimensional constant $C(n)$, the second \approx follows from the spectral gap estimate (2.6).

We now notice that by (3.11), up to possibly increase the value of s_0 , we can apply Theorem 1.1 to each $w(s)$ with $s \geq s_0$: in particular, for every $s \geq s_0$ there exists $\overline{W}(s)$ such that

$$-\Delta \overline{W}(s) = K_0(w(s)) \overline{W}(s)^p, \quad \|\nabla w(s) - \nabla \overline{W}(s)\|_{L^2(\mathbb{R}^n)} \leq C(n) \delta(w(s)). \tag{3.16}$$

By (3.9) and by (3.7),

$$\begin{aligned}
\left| K_0(w(s)) - \frac{1}{1-m} \right|^2 &= \left| \frac{\int_{\mathbb{R}^n} \left(\frac{\Delta w(s)}{w(s)^p} + \frac{1}{1-m} \right) w(s)^{2^*}}{\int_{\mathbb{R}^n} w(s)^{2^*}} \right|^2 \\
&\leq C(n) \int_{\mathbb{R}^n} \left(\frac{\Delta w(s)}{w(s)^p} + \frac{1}{1-m} \right)^2 w(s)^{2^*} = -C(n) \frac{d}{ds} J[w(s)].
\end{aligned} \tag{3.17}$$

Setting $\alpha(s) = [(1-m)K_0(w(s))]^{1/(p-1)}$, the function $\hat{W}(s) = \alpha(s) \overline{W}(s)$ satisfies (3.6), with

$$\|\nabla \hat{W}(s) - \nabla \overline{W}(s)\|_{L^2(\mathbb{R}^n)} \leq C(n) \left| K_0(w(s)) - \frac{1}{1-m} \right|,$$

and hence, by triangle inequality, (3.16), (3.10), and (3.17),

$$\|\nabla w(s) - \nabla \hat{W}(s)\|_{L^2(\mathbb{R}^n)}^2 \leq C(n) \left(\delta(w(s))^2 + \left| K_0(w(s)) - \frac{1}{1-m} \right|^2 \right) \leq -C(n) \frac{d}{ds} J[w(s)].$$

By the minimality property of $W(s)$ and since $J[w(s)]$ and $I[w(s)]$ differ by a constant, we conclude that

$$\|\nabla \rho(s)\|_{L^2(\mathbb{R}^n)}^2 = \|\nabla w(s) - \nabla W(s)\|_{L^2(\mathbb{R}^n)}^2 \leq -C(n) \frac{d}{ds} I[w(s)]$$

for every $s \geq s_0$, and thus, by (3.15),

$$\frac{d}{ds} I[w(s)] \leq -C(n) I[w(s)] \quad \forall s \geq s_0.$$

This proves that, for a constant C_* depending on the initial datum u_0 ,

$$I[w(s)] \leq C_* e^{-\kappa(n)s} \quad \forall s \geq 0 \tag{3.18}$$

(note that the above inequality is trivially true on $[0, s_0]$ by choosing C_* large enough) and hence, applying (3.15) again, we deduce that

$$\int_{\mathbb{R}^n} |\nabla w(s) - \nabla W(s)|^2 \leq C_* e^{-\kappa(n)s} \quad \forall s \geq 0. \tag{3.19}$$

In other words, we have prove that for all s there exists a function $W(s)$ solving (3.6) which is exponentially close to $w(s)$. To conclude the result, we need to show that $W(s)$ is exponentially close to W_∞ . To this aim we notice that, from (3.5),

$$\frac{d}{ds} w^{2^*} = \frac{p+1}{p} w \frac{d}{ds} w^p = \frac{2n}{n+2} \left(\Delta w + \frac{1}{1-m} w^p \right) w,$$

so we obtain

$$|w(s) - w(t)|^{2^*} \leq |w(s)^{2^*} - w(t)^{2^*}| \leq C(n) \left| \int_t^s \left(\frac{\Delta w(r)}{w(r)^p} + \frac{1}{1-m} \right) w(r)^{2^*} dr \right|,$$

and Hölder inequality yields

$$\int_{\mathbb{R}^n} |w(s) - w(t)|^{2^*} \leq C(n) \int_t^s \delta(w(r)) dr \quad \forall s > t > 0. \quad (3.20)$$

Now, by (3.10) and (3.18)

$$\left(\int_k^{k+1} \delta(w(r)) dr \right)^2 \leq \int_k^{k+1} \delta(w(r))^2 \leq C(n) (I[w(k)] - I[w(k+1)]) \leq C_* e^{-\kappa(n)k}$$

which implies

$$\int_t^\infty \delta(w(s)) ds \leq C_* e^{-c(n)t} \quad \forall t > 0.$$

Hence from (3.20) we obtain

$$\int_{\mathbb{R}^n} |w(s) - w(t)|^{2^*} \leq C_* e^{-c(n)t} \quad \forall s > t > 0.$$

From Sobolev inequality and (3.19), it follows that $\{W(s)\}_{s>0}$ is a Cauchy family in $L^{2^*}(\mathbb{R}^n)$ satisfying the exponential bound

$$\int_{\mathbb{R}^n} |W(s) - W(t)|^{2^*} \leq C_* e^{-c(n)t} \quad \forall s > t > 0.$$

Since $\|\nabla W(s) - \nabla W_\infty\|_{L^2(\mathbb{R}^n)} \rightarrow 0$ as $s \rightarrow \infty$ (see (3.12) and (3.19)), we conclude that

$$\int_{\mathbb{R}^n} |W_\infty - W(t)|^{2^*} \leq C_* e^{-c(n)t} \quad \forall t > 0,$$

that combined with (3.19) and Sobolev inequality yields

$$\int_{\mathbb{R}^n} |W_\infty - w(t)|^{2^*} \leq C_* e^{-c(n)t} \quad \forall t > 0. \quad (3.21)$$

Now, to conclude the proof, we argue as follows: let $F : \mathbb{S}^n \rightarrow \mathbb{R}^n$ denote the inverse stereographic projection (see (1.2)), and let $v(s) : \mathbb{S}^n \rightarrow \mathbb{R}$ be defined from $w(s)$ via the transformation

$$w(s, x) = \left(\frac{2}{1 + |x|^2} \right)^{(n-2)/2} v(s, F(x)). \quad (3.22)$$

Then $v(s)$ solves the equation

$$\frac{d}{ds} v^p = \Delta_{\mathbb{S}^n} v + \frac{1}{1-m} v^p - \frac{n(n-2)}{4} v \quad \text{on } (0, \infty) \times \mathbb{S}^n \quad (3.23)$$

(see for instance [Li95] and [dPS01, Equation (2.3)]) and (3.21) translates into

$$\int_{\mathbb{S}^n} |v(s) - v_\infty|^{2^*} \leq C_* e^{-\kappa(n)s} \quad \forall s \geq 0, \quad (3.24)$$

where v_∞ is the stationary solution of (3.23) corresponding to W_∞ under the transformation (3.22). Since $v(s)$ is uniformly bounded away from zero and infinity for s large (see [dPS01, Proposition 5.1]), it follows by (3.24) and parabolic regularity that (up to replacing $\kappa(n)$ by $\kappa(n)/2$)

$$\|v(s) - v_\infty\|_{L^\infty(\mathbb{S}^n)} \leq C_* e^{-\kappa(n)s} \quad \forall s \geq 0.$$

Going back to the original variables, this implies that (3.3) holds, concluding the proof.

BIBLIOGRAPHY

- [AGAP99] A. Ambrosetti, J. Garcia Azorero, and I. Peral. Perturbation of $\Delta u + u^{(N+2)/(N-2)} = 0$, the scalar curvature problem in \mathbf{R}^N , and related topics. *J. Funct. Anal.*, 165(1): 117–149, 1999.
- [AM01] Antonio Ambrosetti and Andrea Malchiodi. On the symmetric scalar curvature problem on S^n . *J. Differential Equations*, 170(1):228–245, 2001.
- [Aub76] Thierry Aubin. Problèmes isopérimétriques et espaces de Sobolev. *J. Differential Geometry*, 11(4):573–598, 1976.
- [Bah96] Abbas Bahri. An invariant for Yamabe-type flows with applications to scalar-curvature problems in high dimension. *Duke Math. J.*, 81(2):323–466, 1996. A celebration of John F. Nash, Jr.
- [BBD⁺09] Adrien Blanchet, Matteo Bonforte, Jean Dolbeault, Gabriele Grillo, and Juan Luis Vázquez. Asymptotics of the fast diffusion equation via entropy estimates. *Arch. Ration. Mech. Anal.*, 191(2):347–385, 2009.
- [BC91] A. Bahri and J.-M. Coron. The scalar-curvature problem on the standard three-dimensional sphere. *J. Funct. Anal.*, 95(1):106–172, 1991.
- [BDGV10] M. Bonforte, J. Dolbeault, G. Grillo, and J. L. Vázquez. Sharp rates of decay of solutions to the nonlinear fast diffusion equation via functional inequalities. *Proc. Natl. Acad. Sci. USA*, 107(38):16459–16464, 2010.
- [BE91] Gabriele Bianchi and Henrik Egnell. A note on the Sobolev inequality. *J. Funct. Anal.*, 100(1):18–24, 1991.
- [BE93] Gabriele Bianchi and Henrik Egnell. A variational approach to the equation $\Delta u + Ku^{(n+2)/(n-2)} = 0$ in \mathbf{R}^n . *Arch. Rational Mech. Anal.*, 122(2):159–182, 1993.
- [BGV10] Matteo Bonforte, Gabriele Grillo, and Juan Luis Vázquez. Special fast diffusion with slow asymptotics: entropy method and flow on a Riemann manifold. *Arch. Ration. Mech. Anal.*, 196(2):631–680, 2010.
- [CENV04] D. Cordero-Erausquin, B. Nazaret, and C. Villani. A mass-transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities. *Adv. Math.*, 182(2):307–332, 2004.
- [Che89] Wen Xiong Chen. Scalar curvatures on S^n . *Math. Ann.*, 283(3):353–365, 1989.
- [Cho92] Bennett Chow. The Yamabe flow on locally conformally flat manifolds with positive Ricci curvature. *Comm. Pure Appl. Math.*, 45(8):1003–1014, 1992.
- [CL97] Wenxiong Chen and Congming Li. A priori estimates for prescribing scalar curvature equations. *Ann. of Math. (2)*, 145(3):547–564, 1997.
- [CY91] Sun-Yung A. Chang and Paul C. Yang. A perturbation result in prescribing scalar curvature on S^n . *Duke Math. J.*, 64(1):27–69, 1991.
- [DKS] Panagiotia Daskalopoulos, Joseph King, and Natasa Sesum. Extinction profile of complete non-compact solutions to the yamabe flow. arxiv preprint 1306.0859.
- [DN85] Wei Yue Ding and Wei-Ming Ni. On the elliptic equation $\Delta u + Ku^{(n+2)/(n-2)} = 0$ and related topics. *Duke Math. J.*, 52(2):485–506, 1985.
- [dPS01] Manuel del Pino and Mariel Sáez. On the extinction profile for solutions of $u_t = \Delta u^{(N-2)/(N+2)}$. *Indiana Univ. Math. J.*, 50(1):611–628, 2001.
- [DS08] Panagiotia Daskalopoulos and Natasa Sesum. On the extinction profile of solutions to fast diffusion. *J. Reine Angew. Math.*, 622:95–119, 2008.
- [ES86] José F. Escobar and Richard M. Schoen. Conformal metrics with prescribed scalar curvature. *Invent. Math.*, 86(2):243–254, 1986.
- [GNN79] B. Gidas, Wei Ming Ni, and L. Nirenberg. Symmetry and related properties via the maximum principle. *Comm. Math. Phys.*, 68(3):209–243, 1979.

- [Heb14] Emmanuel Hebey. *Compactness and stability for nonlinear elliptic equations*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2014. x+291 pp.
- [KW74] Jerry L. Kazdan and F. W. Warner. Curvature functions for compact 2-manifolds. *Ann. of Math. (2)*, 99:14–47, 1974.
- [Li95] Yanyan Li. Prescribing scalar curvature on S^n and related problems. I. *J. Differential Equations*, 120(2):319–410, 1995.
- [Li96] Yanyan Li. Prescribing scalar curvature on S^n and related problems. II. Existence and compactness. *Comm. Pure Appl. Math.*, 49(6):541–597, 1996.
- [LL09] Lishan Lin and Zhaoli Liu. Multi-bump solutions and multi-tower solutions for equations on \mathbb{R}^N . *J. Funct. Anal.*, 257(2):485–505, 2009.
- [LZ15] Man Chun Leung and Feng Zhou. Construction of blow-up sequences for the prescribed scalar curvature equation on S^n . III. Aggregated and towering blow-ups. *Calc. Var. Partial Differential Equations*, 54(3):3009–3035, 2015.
- [Mos73] J. Moser. On a nonlinear problem in differential geometry. In *Dynamical systems (Proc. Sympos., Univ. Bahia, Salvador, 1971)*, pages 273–280. Academic Press, New York, 1973.
- [Ni82] Wei Ming Ni. On the elliptic equation $\Delta u + K(x)u^{(n+2)/(n-2)} = 0$, its generalizations, and applications in geometry. *Indiana Univ. Math. J.*, 31(4):493–529, 1982.
- [Oba72] Morio Obata. The conjectures on conformal transformations of Riemannian manifolds. *J. Differential Geometry*, 6:247–258, 1971/72.
- [Str84] Michael Struwe. A global compactness result for elliptic boundary value problems involving limiting nonlinearities. *Math. Z.*, 187(4):511–517, 1984.
- [Tal76] Giorgio Talenti. Best constant in Sobolev inequality. *Ann. Mat. Pura Appl. (4)*, 110:353–372, 1976.
- [Váz06] Juan Luis Vázquez. *Smoothing and decay estimates for nonlinear diffusion equations*, volume 33 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2006. xiv+234 pp. Equations of porous medium type.
- [Ye94] Rugang Ye. Global existence and convergence of Yamabe flow. *J. Differential Geom.*, 39(1):35–50, 1994.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI PALERMO, VIA ARCHIRAFI 34, 90123 PALERMO, ITALY

E-mail address: giulio.ciraolo@unipa.it

MATHEMATICS DEPARTMENT, ETH ZÜRICH, RÄMISTRASSE 101, CH-8092 ZÜRICH, SWITZERLAND

E-mail address: alessio.figalli@math.ethz.ch

ABDUS SALAM INTERNATIONAL CENTER FOR THEORETICAL PHYSICS, STRADA COSTIERA 11, I-34151, TRIESTE, ITALY. ON LEAVE FROM THE UNIVERSITY OF TEXAS AT AUSTIN

E-mail address: fmaggi@ictp.it